arXiv:1303.1966v1 [quant-ph] 8 Mar 2013

Self-avoiding quantum walks

K. Barr,[1] T. Proctor,[1] B. Hanson,[1] S. Martiel,[1,2] V. Pavlovic,[1,3] A. Bullivant,[1] and V. Kendon[1]

[1]*School of Physics and Astronomy, E. C. Stoner Building, University of Leeds, Leeds LS2 9 JT*
[2]*Lab. I3S, CNRS-UMR 7271, BP 121, 06903 Sophia Antipolis cedex, France*
[3]*Faculty of Science and Mathematics, University of Nis, Serbia*

(Dated: March 11, 2013)

We introduce a new model of the discrete time quantum walk, the self-avoiding quantum walk, which is not allowed to step back onto a position which it has just occupied. This allows us to simulate a dimer and we achieve it by introducing a new type of coin operator. We describe its basic properties and provide numerical evidence that the standard deviation of the walker is constant regardless of the initial state. This contrasts strongly with previously studied coins such as the Grover operator, where the initial condition can be used to control the standard deviation of the walker.

I. INTRODUCTION

The discrete time quantum walk has been extensively studied since its introduction. Initially, various models were proposed [1–3] with a variety of applications in mind. In this paper we are concerned with the transport properties of new type of quantum walk. Quantum walks have been shown to have interesting transport properties in a variety of scenarios. On the line they achieve ballistic transport [4] and they were first shown to have an exponential speedup over the classical random walk on the hypercube by Kempe [5]. The faster hitting times have been used in algorithmic applications [5–7].

The quantum walks introduced thus far concern idealised walkers with no spatial extension. Whilst these have many uses in modelling physical and biological processes [8], in some cases we may want to consider walkers which do have spatial extension, and hence cannot move into space which they are already occupying. Self-avoiding random walks were developed to model precisely such processes, initially the folding of polymers. In some cases, such a restricted model may not be suitable. Continuing the example, polymers can overlap to form loops, but they cannot occupy volume that they already occupy. A type of intermediary between the fully random walk and the self-avoiding walk is known as the non-reversal walk. A non-reversing random walker is allowed to occupy spaces on a lattice that it has previously occupied, but is not allowed to return to the point which it has just come from. In this paper we introduce a quantum version of such a walk, which effectively simulates a dimer. The motivation for studying such a walk is much the same as that for studying the classical version: more realistic simulation of physical systems.

In both the classical and quantum case, the self-avoiding or non-reversal walk on the line is trivial. This is because there are only two degrees of freedom in the movement, so if one of those is prohibited by the model, then unidirectional ballistic transport is obtained. The walks studied in this paper take place over a square lattice, in which case the dynamics are highly non-trivial. The paper proceeds as follows: In Section I A classical self-avoiding walks are introduced in more detail. Then for the sake of comparison the properties of the quantum walk on the square lattice are discussed. The self-avoiding quantum walk is then defined in Section II A and its properties are described in Section II B.

A. Classical self-avoiding random walks

The classical self-avoiding walk has proven difficult to treat analytically, hence the results concerning it have all so far been numerical [9] and there remain many open questions about it. Even enumerating the number of self-avoiding walks has proven very difficult, despite them being so rare that coming upon one by mistake when examining a random walk is highly improbable. If c_n is the number of walks of precisely n steps, then the total number of self-avoiding walks up to length n is $\sum_{n\geq 2} c_n$. Some facts are clear, for example that $c_{n+m} \leq c_m + c_n$. The set of walks of length n concatenated with the set of walks of length m contains not only the self-avoiding walks of length $n+m$ but some which overlap, hence the inequality. However, determining the precise number of walks is difficult, though bounds have been established. The number of self-avoiding walks must be less than the number of non-reversal walks, as these include the self-avoiding walk as a subset. Additionally it is possible to construct subsets of self-avoiding walks which grow as 2^n, so we know that there are between 2^n and 3^n self-avoiding random walks. The best evidence so far suggests that the number of self-avoiding walks of length n is proportional to 2.638^n, and this is provided as a non-rigorous estimate in [10]. The best evidence for this value was obtained by enumerating each such walk of length up to 51 and required a 1024 processor supercomputer [11]. Without new algorithms it is unlikely that we will be able to enumerate much further than this. Guttmann obtained 14,059,415,980,606,050,644,844 walks using this method, however these walks make up less than 1 in 240 million of the possible random walks of this length.

As even counting the walks has proved difficult, it is



FIG. 1. (Color online) Probability distributions arising from a) the DFT coin and b) the Grover coin for the symmetric initial state shown in equation 3

unsurprising that little is known regarding other properties. The property we are most interested in when comparing walks is the standard deviation. The mean squared displacement is conjectured to be $n^{3/2}$ though so far, even a proof that the exponent must be between 1 and 2 is elusive [9].

Another known fact about self-avoiding walks demonstrates a key difference between the self-avoiding walk and its standard and non-reversal counterparts. This is that the self-avoiding walk does not necessarily continue indefinitely. This is because it is possible to reach a lattice site whose only adjacent lattice sites have previously been visited, hence the walker cannot continue. On any given step there is a small probability that the walk will end.

The non-reversal walk is in some ways more tractable. For example, it is clear that on the square lattice there are 3^n such walks, where n is the number of steps traversed by the walker. Its mean squared displacement is $2n$, so it spreads twice as fast as the standard random walk, but slower than the completely self-avoiding walk. There is very little literature on the non-reversal walk, and this tends to examine specific characteristics of the walk relevant to the study of polymer chains [12], rather than its general features.

B. Quantum walks on the square lattice

The properties of the discrete time quantum walk on the square lattice were extensively explored in [13] following initial investigations in [14]. In particular they examined the mean position at time t:

$$\langle r \rangle_t = \sum_{x,y} \sqrt{x^2 + y^2}\, p(x, y, t) \tag{1}$$

so r is the radial distance from the origin, and $p(x, y, t)$ is the probability of finding the walker at position (x, y) at time t. They also characterise the walks in terms of the standard deviation:

$$\sigma = \sqrt{\langle r^2 \rangle - \langle r \rangle^2} \tag{2}$$

The authors use three choices of coin operator. Their first choice behaves like the Hadamard operator for both the ‘left/right’ ($|l\rangle/|r\rangle$) coin states and the ‘up/down’ ($|u\rangle/|d\rangle$) coin states. This operator does not mix the two dimensions, so a two dimensional version of the distribution of the walk on the line is obtained. More interestingly,



FIG. 2. (Color online) Probability distributions arising after 100 steps of a typical self-avoiding quantum walk shown a) over the entire lattice and b) in profile 3

they consider the DFT and Grover operators for a number of initial conditions. These coins are both unbiased, in that they distribute amplitude equally between each coin state. In contrast to the walk on the line, they find that the dynamics of the walk depend strongly on the coin used. Additionally, the dynamics for a specific coin depend strongly on the choice of initial state. The lowest and highest standard deviations obtained for the position of the walker were found using the Grover operator. It was observed that the reason for this is that regardless of the initial state, the distribution forms a central spike, with a ring around it which propagates outwards. The choice of initial condition controls how much amplitude is situated in the central spike, and how much amplitude is situated in the ring. The initial state at the origin for both the distributions plotted in Figure 1 is:

$$
\begin{aligned}
|\psi\rangle &= \frac{1}{2}(|ld\rangle + i|lu\rangle + i|rd\rangle - |ru\rangle) \otimes |0\rangle \\
&= \frac{1}{2}(|l\rangle + i|r\rangle) \otimes (|d\rangle + i|u\rangle) \otimes |0\rangle
\end{aligned}
\tag{3}
$$

The authors of [13] studied all unbiased four dimensional unitary operators with entries equal to either ±1/2 or $\pm i/2$, which when the leading diagonal is selected to be 1/2 gives 640 unitary operators. These operators were found to fall into ten types, with the DFT, Hadamard and Grover all being of different type. Of all operators tested, the Grover operator was found to give rise to both the largest and smallest standard deviation, depending on the initial state.

II. SELF-AVOIDING QUANTUM WALKS

A. Definition

Our definition of a self-avoiding quantum walk is more similar to the classical non-reversal walk than the fully self-avoiding walk. We require that the diagonals of the coin operator are all equal to zero, so that amplitude which has come from a given node cannot return to that node during the next step of the walk. A general coin fulfilling this criteria is specified by:

$$
C^{(SA)} = \begin{pmatrix}
0 & \lambda e^{i\alpha} & \gamma e^{i\beta} & f(\lambda,\gamma)e^{i\theta} \\
-\lambda e^{i(\psi-\phi+\eta)} & 0 & -f(\lambda,\gamma)e^{i(\psi-\theta+\beta)} & \gamma e^{i\psi} \\
\gamma e^{i\eta} & -f(\lambda,\gamma)e^{i(\phi-\theta+\alpha)} & 0 & \lambda e^{i\phi} \\
-f(\lambda,\gamma)e^{i(\theta+\delta+\eta-\phi-\beta)} & -\gamma e^{i(\delta+\alpha-\beta)} & \lambda e^{i\delta} & 0
\end{pmatrix}
\tag{4}
$$

where all of the variables are real and $0 \leq \gamma^2 + \lambda^2 \leq 1$, and $f(\lambda,\gamma) = \sqrt{1-(\lambda^2+\gamma^2)}$. A simple example of such a coin, used to produce the probability distribution shown in Figure 2, takes $\theta = \phi = \pi$ and $\lambda = \gamma = f(\lambda,\gamma) = \frac{1}{\sqrt{3}}$, with all other phase variables being zero. This leads to the following coin:

$$
C^1 = \frac{1}{\sqrt{3}} \begin{pmatrix}
0 & 1 & 1 & -1 \\
1 & 0 & 1 & 1 \\
1 & -1 & 0 & -1 \\
-1 & -1 & 1 & 0
\end{pmatrix}
\tag{5}
$$

We can see that this coin does what it is required to by examining its operation on an arbitrary state

$$
|\psi\rangle = \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix}
\tag{6}
$$

It is clear that

$$
C^1|\psi\rangle = \frac{1}{\sqrt{3}} \begin{pmatrix} b+c-d \\ a+c+d \\ a-b-d \\ c-a-b \end{pmatrix}
\tag{7}
$$

so the initial amplitude in each coin state is redistributed between the other coin states, and none of it is transmitted to the node it came from.

Whilst the walks examined below use real parameters, it is clearly possible to have coins with complex numbers in them, such as

$$
C^2 = \frac{1}{\sqrt{3}} \begin{pmatrix}
0 & i & 1 & i \\
-i & 0 & i & 1 \\
1 & i & 0 & -i \\
i & 1 & i & 0
\end{pmatrix}
\tag{8}
$$

but the properties discussed below have been verified numerically for coins with real entries only.

A self-avoiding walker of length three, which models a trimer, requires a more complex definition. This is because the trimer on a square lattice, presuming that the head and tail are distinguishable, can be in one of three configurations, as shown in Figure 3. These configura-



FIG. 3. The three possible configurations of a self avoiding trimer

tions can then have four possible orientations, facing in the up, down, left and right directions. This means that in order to represent the trimer, a qutrit tensored with a ququat is required, giving rise to a twelve dimensional coin. The head of the trimer can make the same moves as that of the dimer, but now as well as updating the position of the walker, the configuration, which will in general be in a superposition, is also updated.

Whilst it may be possible to define a fully general self-avoiding quantum walk, this will in practise be difficult. Even deciding when a site has been visited has some ambiguity, for instance, can one part of the superposition access sites which another part of the superposition has previously accessed? However, the self-avoiding coin developed here has sufficiently interesting properties for it to be worth exploring regardless, and we now turn to these.

B. Properties

An example of a typical probability distribution arising from a self-avoiding quantum walk is shown in Figure 2. The distribution is shown in 3d and in profile in order to show the similarity of the profile to that of the walk on the line. Regardless of initial condition, the dynamics are similar, tracing out a square with peaks at each corner. The initial condition determines the height and number of distinctive peaks at the corners of the square. The initial condition used to obtain the distribution shown in Figure 2 is the same as that used in Equation 3.

These walks were investigated by varying the parameters in the coin as well as the initial condition. It was found numerically that for all choices of initial condition the mean position (Equation 1) and standard deviation (Equation 2) of the walker is constant at a given time t, as can be seen for the standard deviation in Figure 4. Further investigations suggest that all the moments of the distribution are independent of the initial condition. The parameters which determine the standard deviation are λ and γ, so these values determine some real constant k for each self-avoiding coin so that the standard deviation at time t is equal to kt. This contrasts strongly with standardly used coins, as mentioned in Section I B above, where the initial condition can be used to control the standard deviation of the walk using the Grover coin operator. The mean and standard deviations as a function of time are shown for a variety of coins in Figure 5. Whilst the mean position increases faster than all other coins apart from the Grover, the standard deviation increases more slowly, so in cases where we have freedom to choose our initial condition, the Grover operator is still the best to use if we want to achieve the fastest possible spreading of the amplitude across the lattice.



FIG. 4. Comparison of standard deviation for an arbitrary initial state for (a) the Grover coin and (b) the self-avoiding coin.

III. CONCLUSION

We have introduced a new type of coin for the discrete time quantum walk and shown numerically that it has some notable properties, namely that the mean and standard deviation of the position of the walker is independent of the choice of initial condition. Ideally, an analytic proof that these properties do not depend on the initial condition is required, as well as an analytic expression for the standard deviation in terms of the relevant coin parameters, but these have eluded us thus far. The standard deviation grows linearly with t for much the same reason as it does the walk on the line, as the coin operator always ensures that some amplitude moves away from the origin with each step. However,



FIG. 5. (Color online) Comparison of coin operators in terms of a) mean and b) standard deviation. The initial condition leading to the steepest gradient for each coin was used.

this does not explain why the standard deviation is determined by the choice of λ and γ alone.

A fully self-avoiding quantum walk is yet to be developed and explored, and thus far no continuous time analogues have been proposed. In order to fully capture the folding of polymer chains, the quantum nature of their constituents may need to be accounted for, which would provide a potential application for such a model.

In addition to further investigating the properties of the self-avoiding quantum walk and developing the concept, better comparisons with its classical counterpart may be useful. For example, only walks on square lattices have been considered here, but the self-avoiding random walk has been shown to have macroscopic properties which are independent of the choice of lattice. In order to see if this property carries over into the quantum case, coins of varying dimension are required.

In short, there are many potential avenues through which to explore the self-avoiding quantum walk, and the choices of which direction to take will depend on the desired applications of the walk. The purpose of the work here was simply to find out whether using such a self-avoiding coin operator gives rise to any properties which are not observed using other operators, hopefully provoking further investigation into the uses of such a coin.

ACKNOWLEDGMENTS

KB is funded by the UK Engineering and Physical Sciences Research Council. VK is funded by a Royal Society Research Fellowship. TP, BH, and AB received support from Royal Society summer bursaries and EPSRC.VP was supported by IASTE and SM by JFT Grant ID 15619 and ANR-10-JCJC-0208 CausaQ grant. .

[1] S. Gudder, *Quantum Probability*. Academic Press Inc., CA, USA, 1988.

[2] Aharonov, Y,, Davidovich, L., and Zagury, N., “Quantum random walks,” *Phys. Rev. A*, vol. 2, p. 16871690, 1992.

[3] G. Grossing and A. Zeilinger, “Quantum cellular automata,” *Complex Systems*, vol. 2, p. 197 208, 1988.

[4] A. Ambainis, E. Bach, A. Nayak, A. Vishwanath, and J. Vishwanath, “One-dimensional quantum walks,” *Proc. 33rd Annual ACM STOC*, pp. 60–69, 2001.

[5] J. Kempe, “Quantum random walks hit exponentially faster,” *Probability Th. and Related Fields*, vol. 133 (2), p. 215235., 2005.

[6] A. Childs, R. Cleve, E. Deotto, E. Farhi, S. Gutmann, and D. Spielman, “Exponential algorithmic speedup by quantum walk,” *Proc. 35th annual ACM STOC*, pp. 59–68, 2003.

[7] A. Childs, E. Farhi, and S. Gutmann, “An example of the difference between quantum and classical random walks,” *Quantum Inf. Process.*, vol. 35, no. 1, 2002.

[8] M. Mohseni, P. Rebentrost, S. Lloyd, and A. Aspuru-Guzik, “Environment-assisted quantum walks in photosynthetic energy transfer,” *J. Chem. Phys.*, no. 129, p. 174106, 2008.

[9] B. Hayes, “How to avoid yourself,” *AmSci*, vol. 86, p. 314, 1998.

[10] I. Jensen, “A parallel algorithm for the enumeration of self-avoiding polygons on the square lattice,” *J. Phys. A: Math. Gen*, vol. 36, pp. 5731–5745, 2003.

[11] A. J. Guttmann and A. R. Conway, “Square lattice self-avoiding walks and polygons,” *Ann. Comb.*, vol. 5, pp. 319–345, 2001.

[12] A. Skrilos and S. Chirikjian, “Position and orientation distributions for non-reversal random walks using space-group fourier transforms,” *J Algebr Stat.*, vol. 1, pp. 27–46, 2010.

[13] B. Tregenna, W. Flanagan, R. Maile, and V. Kendon, “Controlling discrete quantum walks: coins and initial states,” *New J. Phys.*, vol. 5, no. 83, 2003.

[14] T. D. Mackay, S. D. Bartlett, L. T. Stephenson, and B. C. Sanders, “Quantum walks in higher dimensions,” *J. Phys. A: Math. Gen.*, vol. 35, 2002.